

Diebold, I.; Schön, U.; Scharf F.; et al. Critical assessment of secondary findings in genes linked to primary arrhythmia syndromes. *Human Mutation*. February 2020:humu.23996. doi:[10.1002/humu.23996](https://doi.org/10.1002/humu.23996)

Diebold, I.; Schön, U.; Horvath, R.; Schwartz, O.; Holinski-Feder, E.; Kolbel, H.; Abicht, A. HADHA and HADHB gene associated phenotypes - Identification of rare variants in a patient cohort by Next Generation Sequencing. *Mol Cell Probes*. 2019;44:14-20. doi:[10.1016/j.mcp.2019.01.003](https://doi.org/10.1016/j.mcp.2019.01.003)

Li, C. G.; Mahon, C.; Sweeney, N. M.; Verschueren, E.; Kantamani, V.; Li, D.; Hennigs, J. K.; Marciano, D. P.; Diebold, I.; Abu-Halawa, O.; et al. PPARgamma Interaction with UBR5/ATMIN Promotes DNA Repair to Maintain Endothelial Homeostasis. *Cell Rep*. 2019;26(5):1333-1343.e7. doi:[10.1016/j.celrep.2019.01.013](https://doi.org/10.1016/j.celrep.2019.01.013)

Abicht, A.; Scharf, F.; Kleinle, S.; Schon, U.; Holinski-Feder, E.; Horvath, R.; Benet-Pages, A.; Diebold, I. Mitochondrial and nuclear disease panel (Mito-aND-Panel): Combined sequencing of mitochondrial and nuclear DNA by a cost-effective and sensitive NGS-based method. *Mol Genet Genomic Med*. 2018;6(6):1188-1198. doi:[10.1002/mgg3.500](https://doi.org/10.1002/mgg3.500)

Trifunov, S.; Pyle, A.; Valentino, M. L.; Liguori, R.; Yu-Wai-Man, P.; Burte, F.; Duff, J.; Kleinle, S.; Diebold, I.; Rugolo, M.; et al. Clonal expansion of mtDNA deletions: different disease models assessed by digital droplet PCR in single muscle cells. *Sci Rep*. 2018;8(1):11682. doi:[10.1038/s41598-018-30143-z](https://doi.org/10.1038/s41598-018-30143-z)

Lamprecht, P.; Humrich, J. Y.; Diebold, I.; Riemekasten, G. Diagnosis of deficiency of adenosine deaminase 2 with early onset polyarteritis nodosa in an adult patient with a novel compound heterozygous CECR1 mutation. *Clin Exp Rheumatol*. 2018;36 Suppl 111(2):177.

Braunisch, M. C.; Gallwitz, H.; Abicht, A.; Diebold, I.; Holinski-Feder, E.; Van Maldergem, L.; Lammens, M.; Kovacs-Nagy, R.; Alhaddad, B.; Strom, T. M.; et al. Extension of the phenotype of biallelic loss-of-function mutations in SLC25A46 to the severe form of pontocerebellar hypoplasia type I. *Clin Genet*. 2018;93(2):255-265. doi:[10.1111/cge.13084](https://doi.org/10.1111/cge.13084)

Hopper, R. K.; Moonen, J.-R. A. J.; Diebold, I.; Cao, A.; Rhodes, C. J.; Tojais, N. F.; Hennigs, J. K.; Gu, M.; Wang, L.; Rabinovitch, M. In Pulmonary Arterial Hypertension, Reduced BMPR2 Promotes Endothelial-to-Mesenchymal Transition via HMGA1 and Its Target Slug. *Circulation*. 2016;133(18):1783-1794. doi:[10.1161/CIRCULATIONAHA.115.020617](https://doi.org/10.1161/CIRCULATIONAHA.115.020617)

Nickel, N. P.; Spiekerkoetter, E.; Gu, M.; Li, C. G.; Li, H.; Kaschwich, M.; Diebold, I.; Hennigs, J. K.; Kim, K.-Y.; Miyagawa, K.; et al. Elafin Reverses Pulmonary Hypertension via Caveolin-1-Dependent Bone Morphogenetic Protein Signaling. *Am J Respir Crit Care Med*. 2015;191(11):1273-1286. doi:[10.1164/rccm.201412-2291OC](https://doi.org/10.1164/rccm.201412-2291OC)

Diebold, I.; Hennigs, J. K.; Miyagawa, K.; Li, C. G.; Nickel, N. P.; Kaschwich, M.; Cao, A.; Wang, L.; Reddy, S.; Chen, P.-I.; et al. BMPR2 preserves mitochondrial function and DNA during reoxygenation to promote endothelial cell survival and reverse pulmonary hypertension. *Cell Metab*. 2015;21(4):596-608. doi:[10.1016/j.cmet.2015.03.010](https://doi.org/10.1016/j.cmet.2015.03.010)

Kleikers, P. W. M.; Wingler, K.; Hermans, J. J. R.; Diebold, I.; Altenhofer, S.; Radermacher, K. A.; Janssen, B.; Gorch, A.; Schmidt, H. H. H. W. NADPH oxidases as a source of oxidative stress and molecular target in ischemia/reperfusion injury. *J Mol Med (Berl)*. 2012;90(12):1391-1406. doi:[10.1007/s00109-012-0963-3](https://doi.org/10.1007/s00109-012-0963-3)

- Diebold, I.; Petry, A.; Sabrane, K.; Djordjevic, T.; Hess, J.; Gorlach, A. The HIF1 target gene NOX2 promotes angiogenesis through urotensin-II. *J Cell Sci.* 2012;125(Pt 4):956-964. doi:[10.1242/jcs.094060](https://doi.org/10.1242/jcs.094060)
- Grunewald, T. G. P.; Diebold, I.; Esposito, I.; Plehm, S.; Hauer, K.; Thiel, U.; da Silva-Buttkus, P.; Neff, F.; Unland, R.; Muller-Tidow, C.; et al. STEAP1 is associated with the invasive and oxidative stress phenotype of Ewing tumors. *Mol Cancer Res.* 2012;10(1):52-65. doi:[10.1158/1541-7786.MCR-11-0524](https://doi.org/10.1158/1541-7786.MCR-11-0524)
- Diebold, I.; Petry, A.; Burger, M.; Hess, J.; Gorlach, A. NOX4 mediates activation of FoxO3a and matrix metalloproteinase-2 expression by urotensin-II. *Mol Biol Cell.* 2011;22(22):4424-4434. doi:[10.1091/mbc.E10-12-0971](https://doi.org/10.1091/mbc.E10-12-0971)
- Su, Y.; Loos, M.; Giese, N.; Hines, O. J.; Diebold, I.; Gorlach, A.; Metzen, E.; Pastorekova, S.; Friess, H.; Buchler, P. PHD3 regulates differentiation, tumour growth and angiogenesis in pancreatic cancer. *Br J Cancer.* 2010;103(10):1571-1579. doi:[10.1038/sj.bjc.6605936](https://doi.org/10.1038/sj.bjc.6605936)
- Diebold, I.; Petry, A.; Djordjevic, T.; Belaiba, R. S.; Fineman, J.; Black, S.; Schreiber, C.; Fratz, S.; Hess, J.; Kietzmann, T.; et al. Reciprocal regulation of Rac1 and PAK-1 by HIF-1alpha: a positive-feedback loop promoting pulmonary vascular remodeling. *Antioxid Redox Signal.* 2010;13(4):399-412. doi:[10.1089/ars.2009.3013](https://doi.org/10.1089/ars.2009.3013)
- Diebold, I.; Flugel, D.; Becht, S.; Belaiba, R. S.; Bonello, S.; Hess, J.; Kietzmann, T.; Gorlach, A. The hypoxia-inducible factor-2alpha is stabilized by oxidative stress involving NOX4. *Antioxid Redox Signal.* 2010;13(4):425-436. doi:[10.1089/ars.2009.3014](https://doi.org/10.1089/ars.2009.3014)
- Diebold, I.; Petry, A.; Hess, J.; Gorlach, A. The NADPH oxidase subunit NOX4 is a new target gene of the hypoxia-inducible factor-1. *Mol Biol Cell.* 2010;21(12):2087-2096. doi:[10.1091/mbc.e09-12-1003](https://doi.org/10.1091/mbc.e09-12-1003)
- Diebold, I.; Djordjevic, T.; Petry, A.; Hatzelmann, A.; Tenor, H.; Hess, J.; Gorlach, A. Phosphodiesterase 2 mediates redox-sensitive endothelial cell proliferation and angiogenesis by thrombin via Rac1 and NADPH oxidase 2. *Circ Res.* 2009;104(10):1169-1177. doi:[10.1161/CIRCRESAHA.109.196592](https://doi.org/10.1161/CIRCRESAHA.109.196592)
- Diebold, I.; Kraicun, D.; Bonello, S.; Gorlach, A. The "PAI-1 paradox" in vascular remodeling. *Thromb Haemost.* 2008;100(6):984-991.
- Diebold, I.; Djordjevic, T.; Hess, J.; Gorlach, A. Rac-1 promotes pulmonary artery smooth muscle cell proliferation by upregulation of plasminogen activator inhibitor-1: role of NFkappaB-dependent hypoxia-inducible factor-1alpha transcription. *Thromb Haemost.* 2008;100(6):1021-1028.
- Herkert, O.; Diebold, I.; Brandes, R. P.; Hess, J.; Busse, R.; Gorlach, A. NADPH oxidase mediates tissue factor-dependent surface procoagulant activity by thrombin in human vascular smooth muscle cells. *Circulation.* 2002;105(17):2030-2036. doi:[10.1161/01.cir.0000014611.28864.1e](https://doi.org/10.1161/01.cir.0000014611.28864.1e)
- Gorlach, A.; Diebold, I.; Schini-Kerth, V. B.; Berchner-Pfannschmidt, U.; Roth, U.; Brandes, R. P.; Kietzmann, T.; Busse, R. Thrombin activates the hypoxia-inducible factor-1 signaling pathway in vascular smooth muscle cells: Role of the p22(phox)-containing NADPH oxidase. *Circ Res.* 2001;89(1):47-54. doi:[10.1161/hh1301.092678](https://doi.org/10.1161/hh1301.092678)